

6.79

**6.79** An incompressible, viscous fluid is placed between horizontal, infinite, parallel plates as is shown in Fig. P6.94. The two plates move in opposite directions with constant velocities,  $U_1$  and  $U_2$ , as shown. The pressure gradient in the  $x$  direction is zero and the only body force is due to the fluid weight. Use the Navier-Stokes equations to derive an expression for the velocity distribution between the plates. Assume laminar flow.




FIGURE P6.94

For the specified conditions,  $v=0$ ,  $w=0$ ,  $\frac{\partial p}{\partial x}=0$ , and  $g_x=0$ , so that the  $x$ -component of the Navier-Stokes equations (Eq. 6.127a) reduces to

$$\frac{d^2 u}{dy^2} = 0 \quad (1)$$

Integration of Eq. (1) yields

$$u = C_1 y + C_2 \quad (2)$$

For  $y=0$ ,  $u = -U_2$  and therefore from Eq. (2)

$$C_2 = -U_2$$

For  $y=b$ ,  $u = U_1$ , so that

$$U_1 = C_1 b - U_2$$

or

$$C_1 = \frac{U_1 + U_2}{b}$$

Thus,

$$\underline{\underline{u = \left( \frac{U_1 + U_2}{b} \right) y - U_2}}$$